

# Introduction

---

Lijun Zhang

[zlj@nju.edu.cn](mailto:zlj@nju.edu.cn)

<http://cs.nju.edu.cn/zlj>





# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Mathematical Optimization (1)

## □ Optimization Problem

$$\begin{aligned} \min \quad & f_0(x) \\ \text{s. t.} \quad & f_i(x) \leq b_i, \quad i = 1, \dots, m \end{aligned}$$

- Optimization Variable:  $x = (x_1, \dots, x_n)$
- Objective Function:  $f_0: \mathbf{R}^n \rightarrow \mathbf{R}$
- Constraint Functions:  $f_i: \mathbf{R}^n \rightarrow \mathbf{R}$

## □ $x^*$ is called optimal or a solution

- $f_i(x^*) \leq b_i, i = 1, \dots, m$
- For any  $z$  with  $f_i(z) \leq b_i$ , we have  $f_0(z) \geq f_0(x^*)$



# Mathematical Optimization (2)

---

## □ Linear Problem

$$f_i(\alpha x + \beta y) = \alpha f_i(x) + \beta f_i(y)$$

- for all  $x, y \in \mathbf{R}^n$  and all  $\alpha, \beta \in \mathbf{R}$

## □ Nonlinear Program

- If the optimization problem is not linear

## □ Convex Optimization Problem

$$f_i(\alpha x + \beta y) \leq \alpha f_i(x) + \beta f_i(y)$$

- for all  $x, y \in \mathbf{R}^n$  and all  $\alpha, \beta \in \mathbf{R}$  with  $\alpha + \beta = 1, \alpha \geq 0, \beta \geq 0$



# Applications

---

$$\begin{array}{ll} \min & f_0(x) \\ \text{s. t.} & f_i(x) \leq b_i, \quad i = 1, \dots, m \end{array}$$

## □ Abstraction

- $x$  represents the choice made
- $f_i(x) \leq b_i$  represent firm requirements that limit the possible choices
- $f_0(x)$  represents the cost of choosing  $x$
- A solution corresponds to a choice that has minimum cost, among all choices that meet the requirements



# Portfolio Optimization

---

## □ Variables

- $x_i$  represents the investment in the  $i$ -th asset
- $x \in \mathbf{R}^n$  describes the overall portfolio allocation across the set of asset

## □ Constraints

- A limit on the budget the requirement
- Investments are nonnegative
- A minimum acceptable value of expected return for the whole portfolio

## □ Objective

- Minimize the variance of the portfolio return



# Device Sizing

---

## □ Variables

- $x \in \mathbf{R}^n$  describes the widths and lengths of the devices

## □ Constraints

- Limits on the device sizes
- Timing requirements
- A limit on the total area of the circuit

## □ Objective

- Minimize the total power consumed by the circuit





# Data Fitting

---

## □ Variables

- $x \in \mathbf{R}^n$  describes parameters in the model

## □ Constraints

- Prior information
- Required limits on the parameters (such as nonnegativity)

## □ Objective

- Minimize the prediction error between the observed data and the values predicted by the model



# Solving Optimization Problems

---

## □ General Optimization Problem

- Very difficult to solve
- Constraints can be very complicated, the number of variables can be very large
- Methods involve some compromise, e.g., computation time, or suboptimal solution

## □ Exceptions

- Least-squares problems
- Linear programming problems
- Convex optimization problems



# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Least-squares Problems (1)

## □ The Problem

$$\min \|Ax - b\|_2^2 = \sum_{i=1}^k (a_i^\top x - b_i)^2$$

- $A \in \mathbf{R}^{k \times n}$ ,  $a_i^\top$  is the  $i$ -th row of  $A$ ,  $b \in \mathbf{R}^k$
- $x \in \mathbf{R}^n$  is the optimization variable



How to solve it?



# Least-squares Problems (1)

---

## □ The Problem

$$\min \|Ax - b\|_2^2 = \sum_{i=1}^k (a_i^\top x - b_i)^2$$

- $A \in \mathbf{R}^{k \times n}$ ,  $a_i^\top$  is the  $i$ -th row of  $A$ ,  $b \in \mathbf{R}^k$
- $x \in \mathbf{R}^n$  is the optimization variable

## □ Setting the gradient to be 0

$$\begin{aligned} & 2A^\top (Ax - b) = 0 \\ \Rightarrow & A^\top Ax = A^\top b \\ \Rightarrow & x = (A^\top A)^{-1} A^\top b \end{aligned}$$



# Least-squares Problems (2)

---

## □ A Set of Linear Equations

$$A^T A x = A^T b$$

## □ Solving least-squares problems

- Reliable and efficient algorithms and software
- Computation time proportional to  $n^2 k$  ( $A \in \mathbf{R}^{k \times n}$ ); less if structured
- A mature technology
- Challenging for **extremely large** problems



# Using Least-squares

---

- Easy to Recognize
- Weighted least-squares

$$\sum_{i=1}^k w_i (a_i^\top x - b_i)^2$$

- Different importance



# Using Least-squares

- Easy to Recognize
- Weighted least-squares

$$\sum_{i=1}^k w_i (a_i^\top x - b_i)^2 = \sum_{i=1}^k (\sqrt{w_i} a_i^\top x - \sqrt{w_i} b_i)^2$$

- Different importance

- Regularization

$$\sum_{i=1}^k (a_i^\top x - b_i)^2 + \rho \sum_{i=1}^n x_i^2$$

- More stable





# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Linear Programming

---

## □ The Problem

$$\begin{aligned} \min \quad & c^T x \\ \text{s. t.} \quad & a_i^T x \leq b_i, \quad i = 1, \dots, m \end{aligned}$$

- $c, a_1, \dots, a_m \in \mathbf{R}^n, b_1, \dots, b_m \in \mathbf{R}$

## □ Solving Linear Programs

- No analytical formula for solution
- Reliable and efficient algorithms and software
- Computation time proportional to  $n^2m$  if  $m \geq n$ ; less with structure
- A mature technology
- Challenging for **extremely large** problems



# Using Linear Programming

- Not as easy to recognize
- Chebyshev Approximation Problem

$$\min \max_{i=1, \dots, k} |a_i^\top x - b_i|$$

$$\begin{aligned} & \iff \min t \\ & \text{s. t. } t = \max_{i=1, \dots, k} |a_i^\top x - b_i| \end{aligned}$$

$$\begin{aligned} & \iff \min t \\ & \text{s. t. } t \geq |a_i^\top x - b_i|, i = 1, \dots, k \end{aligned}$$

$$\begin{aligned} & \iff \min t \\ & \text{s. t. } -t \leq a_i^\top x - b_i \leq t, i = 1, \dots, k \end{aligned}$$



# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary




# Convex Optimization


## □ Why Convexity?

“ The great watershed in optimization isn't between linearity and nonlinearity, but convexity and nonconvexity.”

— R. Rockafellar, SIAM Review 1993



**Non-Convex Optimization**



**Convex Optimization**




# Convex Optimization


## □ Why Convexity?

“ The great watershed in optimization  
nonlinearity, but convexity and no  
— R. ... 1993

Local minimizers  
are also global  
minimizers.



Non-Convex Optimization



Convex Optimization



# Convex Optimization Problems (1)

---

## □ The Problem

$$\begin{array}{ll} \min & f_0(x) \\ \text{s. t.} & f_i(x) \leq b_i, \quad i = 1, \dots, m \end{array}$$

- Functions  $f_0, \dots, f_m: \mathbf{R}^n \rightarrow \mathbf{R}$  are convex:

$$f_i(\alpha x + \beta y) \leq \alpha f_i(x) + \beta f_i(y)$$

for all  $x, y \in \mathbf{R}^n$  and all  $\alpha, \beta \in \mathbf{R}$  with  $\alpha + \beta = 1$ ,  $\alpha \geq 0$ ,  $\beta \geq 0$

- Least-squares and linear programs as special cases



## Convex Optimization Problems (2)

---

### □ Solving Convex Optimization Problems

- No analytical solution
- Reliable and efficient algorithms (e.g., interior-point methods)
- Computation time (roughly) proportional to  $\max\{n^3, n^2m, F\}$ 
  - ✓  $F$  is cost of evaluating  $f_i'$ s and their first and second derivatives
- **Almost** a technology





# Using Convex Optimization


---

- Often difficult to recognize
- Many tricks for transforming problems into convex form
- Surprisingly many problems can be solved via convex optimization



# An Example (1)

□  $m$  lamps illuminating  $n$  patches



■ Intensity  $I_k$  at patch  $k$  depends linearly on lamp powers  $p_j$

$$I_k = \sum_{j=1}^m a_{kj} p_j, \quad a_{kj} = r_{kj}^{-2} \max\{\cos\theta_{kj}, 0\}$$



## An Example (2)

---

- Achieve desired illumination  $I_{\text{des}}$  with bounded lamp powers

$$\begin{aligned} \min \quad & \max_{k=1,\dots,n} |\log I_k - \log I_{\text{des}}| \\ \text{s. t.} \quad & 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$



How to solve it?



## An Example (3)

1. Use uniform power:  $p_j = p$ , vary  $p$
2. Use least-squares

$$\min \sum_{i=1}^k (I_k - I_{\text{des}})^2 = \sum_{i=1}^k \left( \sum_{j=1}^m a_{kj} p_j - I_{\text{des}} \right)^2$$

- Round  $p_j$  if  $p_j > p_{\text{max}}$  or  $p_j < 0$

3. Use weighted least-squares

$$\min \sum_{i=1}^k (I_k - I_{\text{des}})^2 + \sum_{j=1}^m w_j \left( p_j - \frac{p_{\text{max}}}{2} \right)^2$$

- Adjust weights  $w_j$  until  $0 \leq p_j \leq p_{\text{max}}$



## An Example (4)

---

### 4. Use linear programming

$$\begin{aligned} \min \quad & \max_{k=1,\dots,n} |I_k - I_{\text{des}}| \\ \text{s. t.} \quad & 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$

### 5. Use convex optimization

$$\begin{aligned} \min \quad & \max_{k=1,\dots,n} |\log I_k - \log I_{\text{des}}| \\ \text{s. t.} \quad & 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \min \quad & \max_{k=1,\dots,n} \max \left( \log \frac{I_k}{I_{\text{des}}}, \log \frac{I_{\text{des}}}{I_k} \right) \\ \text{s. t.} \quad & 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$




# An Example (5)

$$\begin{aligned} \Leftrightarrow \quad & \min \quad \max_{k=1,\dots,n} \max \left( \frac{I_k}{I_{\text{des}}}, \frac{I_{\text{des}}}{I_k} \right) \\ & \text{s. t.} \quad 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \quad & \min \quad \max_{k=1,\dots,n} h \left( \frac{I_k}{I_{\text{des}}} \right) \\ & \text{s. t.} \quad 0 \leq p_j \leq p_{\text{max}}, j = 1, \dots, m \end{aligned}$$

■  $h(u) = \max \left( u, \frac{1}{u} \right)$





# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Nonlinear Optimization

---

- An optimization problem when the objective or constraint functions are not linear, but not known to be convex
  
- Sadly, there are no effective methods for solving the general nonlinear programming problem
  - Could be NP-hard
  
- We need **compromise**





# Local Optimization Methods

---

- Find a point that minimizes  $f_0$  among feasible points near it
  - The compromise is to give up seeking the optimal  $x$
- Fast, can handle large problems
  - Differentiability
- Require initial guess
- Provide no information about distance to (global) optimum
- Local optimization methods are **more art than technology**



# Comparisons

---

	Problem Formulation	Solving the Problem
Local Optimization Methods for Nonlinear Programming	Straightforward	<b>Art</b>
Convex Optimization	<b>Art</b>	Standard



# Global Optimization

---

- Find the global solution
  - The compromise is efficiency
- Worst-case complexity grows exponentially with problem size
  
- Worst-case Analysis
  - Whether the worst-case value is acceptable
  - A local optimization method can be tried

# Role of Convex Optimization in Nonconvex Problems

---



- Initialization for local optimization
  - An approximate, but convex, formulation
  
- Convex heuristics for nonconvex optimization
  - Sparse solutions (compressive sensing)
  
- Bounds for global optimization
  - Relaxation
  - Lagrangian relaxation



# Outline

---

- Mathematical Optimization
- Least-squares
- Linear Programming
- Convex Optimization
- Nonlinear Optimization
- Summary



# Summary

---

- Mathematical Optimization
- Least-squares
  - Closed-form Solution
- Linear Programming
  - Efficient algorithms
- Convex Optimization
  - Efficient algorithms, Modeling is an art
- Nonlinear Optimization
  - Compromises, Optimization is an Art