

# Mathematical Background

---

Lijun Zhang

[zlj@nju.edu.cn](mailto:zlj@nju.edu.cn)

<http://cs.nju.edu.cn/zlj>





# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra



# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra



# Inner product

---

- Inner product on  $\mathbf{R}^n$

$$\langle x, y \rangle = x^T y = \sum_{i=1}^n x_i y_i, \quad x, y \in \mathbf{R}^n$$

- Euclidean norm, or  $l_2$ -norm

$$\|x\|_2 = (x^T x)^{1/2} = (x_1^2 + \cdots + x_n^2)^{1/2}, \quad x \in \mathbf{R}^n$$

- Cauchy-Schwarz inequality

$$|x^T y| \leq \|x\|_2 \|y\|_2, \quad x, y \in \mathbf{R}^n$$

- Angle between nonzero vectors  $x, y \in \mathbf{R}^n$

$$\angle(x, y) = \cos^{-1} \left( \frac{x^T y}{\|x\|_2 \|y\|_2} \right), \quad x, y \in \mathbf{R}^n$$



# Inner product

- Inner product on  $\mathbf{R}^{m \times n}$ ,  $X, Y \in \mathbf{R}^{m \times n}$

$$\langle X, Y \rangle = \text{tr}(X^T Y) = \sum_{i=1}^m \sum_{j=1}^n X_{ij} Y_{ij}$$

Here  $\text{tr}(\cdot)$  denotes trace of a matrix

- Frobenius norm of a matrix  $X \in \mathbf{R}^{m \times n}$

$$\|X\|_F = (\text{tr}(X^T X))^{1/2} = \left( \sum_{i=1}^m \sum_{j=1}^n X_{ij}^2 \right)^{1/2}$$

- Inner product on  $\mathbf{S}^n$

$$\langle X, Y \rangle = \text{tr}(XY) = \sum_{i=1}^n \sum_{j=1}^n X_{ij} Y_{ij} = \sum_{i=1}^n X_{ii} Y_{ii} + 2 \sum_{i < j} X_{ij} Y_{ij}$$



# Norms

---

- A function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$  with  $\text{dom } f = \mathbf{R}^n$  is called a norm if
  - $f$  is nonnegative:  $f(x) \geq 0$  for all  $x \in \mathbf{R}^n$
  - $f$  is definite:  $f(x) = 0$  only if  $x = 0$
  - $f$  is homogeneous:  $f(tx) = |t|f(x)$ , for all  $x \in \mathbf{R}^n$  and  $t \in \mathbf{R}$
  - $f$  satisfies the triangle inequality:  
 $f(x + y) \leq f(x) + f(y)$ , for all  $x, y \in \mathbf{R}^n$
- Distance
  - Between vectors  $x$  and  $y$  as the length of their difference, i.e.,  
$$\text{dist}(x, y) = \|x - y\|$$



# Norms

---

## □ Unit ball

- The set of all vectors with norm less than or equal to one,

$$\mathcal{B} = \{x \in \mathbf{R}^n \mid \|x\| \leq 1\}$$

is called the unit ball of the norm  $\|\cdot\|$ .

- The unit ball satisfies the following properties:
  - ✓  $\mathcal{B}$  is symmetric about the origin, i.e.,  $x \in \mathcal{B}$  if and only if  $-x \in \mathcal{B}$
  - ✓  $\mathcal{B}$  is convex
  - ✓  $\mathcal{B}$  is closed, bounded, and has nonempty interior
- Conversely, if  $C \subseteq \mathbf{R}^n$  is any set satisfying these three conditions, then it is the unit ball of a norm:

$$\|x\| = (\sup\{t \geq 0 \mid tx \in C\})^{-1}$$



# Norms

---

## □ Some common norms on $\mathbf{R}^n$

- Sum-absolute-value, or  $l_1$ -norm

$$\|x\|_1 = |x_1| + \cdots + |x_n|, x \in \mathbf{R}^n$$

- Chebyshev or  $l_\infty$ -norm

$$\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$$

- $l_p$ -norm,  $p \geq 1$

$$\|x\|_p = (|x_1|^p + \cdots + |x_n|^p)^{1/p}$$

- For  $P \in \mathbf{S}_{++}^n$ ,  $P$ -quadratic norm is

$$\|x\|_P = (x^\top P x)^{1/2} = \|P^{1/2} x\|_2$$





# Norms

---

□ Some common norms on  $\mathbf{R}^{m \times n}$

■ Sum-absolute-value norm

$$\|X\|_{\text{sav}} = \sum_{i=1}^m \sum_{j=1}^n |X_{ij}|$$

■ Maximum-absolute-value norm

$$\|X\|_{\text{mav}} = \max\{|X_{ij}| \mid i = 1, \dots, m, j = 1, \dots, n\}$$



# Norms

---

## □ Equivalence of norms

- Suppose that  $\|\cdot\|_a$  and  $\|\cdot\|_b$  are norms on  $\mathbf{R}^n$ , there exist positive constants  $\alpha$  and  $\beta$ , for all  $x \in \mathbf{R}^n$

$$\alpha \|x\|_a \leq \|x\|_b \leq \beta \|x\|_a$$

- If  $\|\cdot\|$  is any norm on  $\mathbf{R}^n$ , then there exists a quadratic norm  $\|\cdot\|_P$  for which

$$\|x\|_P \leq \|x\| \leq \sqrt{n} \|x\|_P$$

holds for all  $x$



# Norms

## □ Operator norms

- Suppose  $\|\cdot\|_a$  and  $\|\cdot\|_b$  are norms on  $\mathbf{R}^m$  and  $\mathbf{R}^n$ , respectively. Operator norm of  $X \in \mathbf{R}^{m \times n}$  induced by  $\|\cdot\|_a$  and  $\|\cdot\|_b$  is

$$\|X\|_{a,b} = \sup\{\|Xu\|_a \mid \|u\|_b \leq 1\}$$

- When  $\|\cdot\|_a$  and  $\|\cdot\|_b$  are Euclidean norms, the operator norm of  $X$  is its maximum singular value, and is denoted  $\|X\|_2$

$$\|X\|_2 = \sigma_{\max}(X) = (\lambda_{\max}(X^T X))^{1/2}$$

- ✓ Spectral norm or  $\ell_2$ -norm



# Norms

## □ Operator norms

- The norm induced by the  $\ell_\infty$ -norm on  $\mathbf{R}^m$  and  $\mathbf{R}^n$ , denoted  $\|X\|_\infty$ , is the max-row-sum norm,

$$\|X\|_\infty = \sup\{\|Xu\|_\infty \mid \|u\|_\infty \leq 1\} = \max_{i=1,\dots,m} \sum_{j=1}^n |X_{ij}|$$

- The norm induced by the  $\ell_1$ -norm on  $\mathbf{R}^m$  and  $\mathbf{R}^n$ , denoted  $\|X\|_1$ , is the max-column-sum norm,

$$\|X\|_1 = \max_{j=1,\dots,n} \sum_{i=1}^m |X_{ij}|$$



# Norms

## □ Dual norm

- Let  $\|\cdot\|$  be a norm on  $\mathbf{R}^n$
- The associated dual norm, denoted  $\|\cdot\|_*$ , is defined as

$$\|z\|_* = \sup\{z^\top x \mid \|x\| \leq 1\}$$

- We have the **inequality**

$$z^\top x \leq \|x\| \|z\|_*$$

$$z^\top x = z^\top \frac{x}{\|x\|} \cdot \|x\| \leq \|z\|_* \|x\|$$

$$z^\top \frac{x}{\|x\|} \leq \sup\{z^\top x \mid \|x\| \leq 1\} = \|z\|_*$$



# Norms

## □ Dual norm

- Let  $\|\cdot\|$  be a norm on  $\mathbf{R}^n$
- The associated dual norm, denoted  $\|\cdot\|_*$ , is defined as

$$\|z\|_* = \sup\{z^T x \mid \|x\| \leq 1\}$$

- We have the **inequality**

$$z^T x \leq \|x\| \|z\|_*$$

- The dual of Euclidean norm

$$\sup\{z^T x \mid \|x\|_2 \leq 1\} = \|z\|_2$$

- The dual of the  $\ell_\infty$ -norm

$$\sup\{z^T x \mid \|x\|_\infty \leq 1\} = \|z\|_1$$



# Norms

---

## □ Dual norm

- Let  $\|\cdot\|$  be a norm on  $\mathbf{R}^n$
- The associated dual norm, denoted  $\|\cdot\|_*$ , is defined as

$$\|z\|_* = \sup\{z^\top x \mid \|x\| \leq 1\}$$

- We have the **inequality**

$$z^\top x \leq \|x\| \|z\|_*$$

- The dual of the dual norm

$$\|\cdot\|_{**} = \|\cdot\|$$



# Norms

---

## □ Dual Norm

- The dual of  $\ell_p$ -norm is the  $\ell_q$ -norm such that

$$\frac{1}{p} + \frac{1}{q} = 1$$

- The dual of the  $\ell_2$ -norm on  $\mathbf{R}^{m \times n}$  is the **nuclear norm**

$$\begin{aligned}\|Z\|_{2^*} &= \sup\{\text{tr}(Z^T X) \mid \|X\|_2 \leq 1\} \\ &= \sigma_1(Z) + \cdots + \sigma_r(Z) = \text{tr}[(Z^T Z)^{1/2}]\end{aligned}$$





# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra



# Analysis

---

## □ Interior and Open Set

- An element  $x \in C \subseteq \mathbf{R}^n$  is called an interior point of  $C$  if there exists an  $\epsilon > 0$  for which

$$\{y \mid \|y - x\|_2 \leq \epsilon\} \subseteq C$$

i.e., there exists a ball centered at  $x$  that lies entirely in  $C$

- The set of all points interior to  $C$  is called the interior of  $C$  and is denoted  $\text{int } C$
- A set  $C$  is open if  $\text{int } C = C$



# Analysis

---

## □ Closed Set and Boundary

- A set  $C \subseteq \mathbf{R}^n$  is closed if its complement is open

$$\mathbf{R}^n \setminus C = \{x \in \mathbf{R}^n \mid x \notin C\}$$

- The closure of a set  $C$  is defined as

$$\text{cl } C = \mathbf{R}^n \setminus \text{int}(\mathbf{R}^n \setminus C)$$

- The boundary of the set  $C$  is defined as

$$\text{bd } C = \text{cl } C \setminus \text{int } C$$

- ✓  $C$  is closed if it contains its boundary. It is open if it contains no boundary points



# Analysis

---

## □ Supremum and infimum

- The least upper bound or supremum of the set  $C$  is denoted  $\sup C$
- The greatest lower bound or infimum of the set  $C$  is denoted  $\inf C$

$$\inf C = -(\sup -C)$$



# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra



# Functions

---

## □ Notation

$$f: A \rightarrow B$$

- $\text{dom } f \subseteq A$

## □ An example $f: \mathbf{S}^n \rightarrow \mathbf{R}$

$$f(X) = \log \det X$$

- $\text{dom } f = \mathbf{S}_{++}^n$



# Functions

## □ Continuity

- A function  $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$  is continuous at  $x \in \text{dom } f$  if for all  $\epsilon > 0$  there exists a  $\delta$  such that  $y \in \text{dom } f, \|y - x\|_2 \leq \delta \Rightarrow \|f(y) - f(x)\|_2 \leq \epsilon$
- $f$  is continuous if it is continuous at every point

## □ Closed functions

- A function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$  is closed if, for each  $\alpha \in \mathbf{R}$ , the sublevel set

$$\{x \in \text{dom } f \mid f(x) \leq \alpha\}$$

is closed. This is equivalent to

$\text{epi } f = \{(x, t) \in \mathbf{R}^{n+1} \mid x \in \text{dom } f, f(x) \leq t\}$  is closed



# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra





# Derivatives

## □ Definition

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $x \in \text{int dom } f$ . The function  $f$  is differentiable at  $x$  if there exists a matrix  $Df(x) \in \mathbf{R}^{m \times n}$  that satisfies

$$\lim_{z \in \text{dom } f, z \neq x, z \rightarrow x} \frac{\|f(z) - f(x) - Df(x)(z - x)\|_2}{\|z - x\|_2} = 0$$

in which case we refer to  $Df(x)$  as the derivative (or Jacobian) of  $f$  at  $x$

- $f$  is differentiable if  $\text{dom } f$  is open, and it is differentiable at every point



# Derivatives

---

## □ Definition

- The affine function of  $z$  given by

$$f(x) + Df(x)(z - x)$$

is called the **first-order approximation** of  $f$  at (or near)  $x$

$$Df(x)_{ij} = \frac{\partial f_i(x)}{\partial x_j}, i = 1, \dots, m, j = 1, \dots, n$$



# Derivatives

## □ Gradient

- When  $f$  is real-valued (i.e.,  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ ) the derivative  $Df(x)$  is a  $1 \times n$  matrix (it is a row vector). Its transpose is called the gradient of the function:

$$\nabla f(x) = Df(x)^\top$$

which is a column vector (in  $\mathbf{R}^n$ ). Its components are the partial derivatives of  $f$ :

$$\nabla f(x)_i = \frac{\partial f(x)}{\partial x_i}, i = 1, \dots, n$$

- The **first-order approximation** of  $f$  at a point  $x \in \text{int dom } f$  can be expressed as (the affine function of  $z$ )

$$f(x) + \nabla f(x)^\top (z - x)$$



# Derivatives

---

## □ Examples

$$f(x) = \frac{1}{2}x^\top Px + q^\top x + r, \quad P \in \mathbf{S}^n$$

$$\nabla f(x) = Px + q$$

$$f(X) = \log \det X, \text{ dom } f = \mathbf{S}_{++}^n$$

$$\nabla f(X) = X^{-1}$$



# Derivatives

---

## □ Chain rule

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$  is differentiable at  $x \in \text{int dom } f$  and  $g: \mathbf{R}^m \rightarrow \mathbf{R}^p$  is differentiable at  $f(x) \in \text{int dom } g$ .

Define the composition  $h: \mathbf{R}^n \rightarrow \mathbf{R}^p$  by  $h(z) = g(f(z))$ . Then  $h$  is differentiable at  $x$ , with derivative

$$Dh(x) = Dg(f(x))Df(x)$$

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g: \mathbf{R} \rightarrow \mathbf{R}$ , and  $h(x) = g(f(x))$

$$\nabla h(x) = g'(f(x))\nabla f(x)$$



# Derivatives

---

## □ Composition of Affine Function

$$g(x) = f(Ax + b)$$

$$\nabla g(x) = A^\top \nabla f(Ax + b)$$

$$f: \mathbf{R}^n \rightarrow \mathbf{R}, \quad g: \mathbf{R} \rightarrow \mathbf{R}$$

$$g(t) = f(x + tv), \quad x, v \in \mathbf{R}^n$$

$$g'(t) = v^\top \nabla f(x + tv)$$



# Example 1

---

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$g(y) = \log \sum_{i=1}^m \exp(y_i)$$

$$\nabla g(y) = \frac{1}{\sum_{i=1}^m \exp y_i} \begin{bmatrix} \exp y_1 \\ \vdots \\ \exp y_m \end{bmatrix}$$



# Example 1

---

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$\nabla f(x) = A^\top \nabla g(Ax + b)$$

$$\nabla g(y) = \frac{1}{\sum_{i=1}^m \exp y_i} \begin{bmatrix} \exp y_1 \\ \vdots \\ \exp y_m \end{bmatrix}$$





# Example 1

---

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$\nabla f(x) = A^\top \nabla g(Ax + b) = \frac{1}{\mathbf{1}^\top z} A^\top z$$

$$z = \begin{bmatrix} \exp a_1^\top x + b_1 \\ \vdots \\ \exp a_m^\top x + b_m \end{bmatrix}$$



## Example 2

□ Consider the function

$$f(x) = \log \det(F_0 + x_1 F_1 + \cdots + x_n F_n)$$

■ where  $F_0, \dots, F_n \in \mathbf{S}^p$

□  $f(x) = g(F_0 + x_1 F_1 + \cdots + x_n F_n)$

$$g(X) = \log \det X$$

$$\frac{\partial f(x)}{\partial x_i} = \text{tr}(F_i \nabla \log \det(F)) = \text{tr}(F^{-1} F_i)$$



$$g(t) = f(x + tv), \quad x, v \in \mathbf{R}^n$$

$$g'(t) = v^\top \nabla f(x + tv)$$



## Example 2

---

□ Consider the function

$$f(x) = \log \det(F_0 + x_1 F_1 + \cdots + x_n F_n)$$

■ where  $F_0, \dots, F_n \in \mathbf{S}^p$

□  $f(x) = g(F_0 + x_1 F_1 + \cdots + x_n F_n)$

$$g(X) = \log \det X$$

$$\frac{\partial f(x)}{\partial x_i} = \text{tr}(F_i \nabla \log \det(F)) = \text{tr}(F^{-1} F_i)$$

$$\nabla f(x) = \begin{bmatrix} \text{tr}(F^{-1} F_1) \\ \vdots \\ \text{tr}(F^{-1} F_n) \end{bmatrix}$$



# Second Derivative

## □ Definition

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ . The second derivative or Hessian matrix of  $f$  at  $x \in \text{int dom } f$ , denoted  $\nabla^2 f(x)$ , is given by

$$\nabla^2 f(x)_{ij} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}, i = 1, \dots, n, j = 1, \dots, n.$$

## □ Second-order Approximation

$$f(x) + \nabla f(x)^\top (z - x) + \frac{1}{2} (z - x)^\top \nabla^2 f(x) (z - x)$$



# Derivatives

## □ Examples

$$f(x) = \frac{1}{2}x^\top Px + q^\top x + r, \quad P \in \mathbf{S}^n$$

$$\nabla f(x) = Px + q$$

$$\nabla^2 f(x) = P$$

$$f(X) = \log \det X, \text{ dom } f = \mathbf{S}_{++}^n$$

$$\nabla f(X) = X^{-1}$$

$$f(X) + \text{tr}(X^{-1}(Z - X)) - \frac{1}{2}\text{tr}(X^{-1}(Z - X)X^{-1}(Z - X))$$



# Second Derivative

## □ Chain rule

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g: \mathbf{R} \rightarrow \mathbf{R}$ , and  $h(x) = g(f(x))$

$$\nabla h(x) = g'(f(x))\nabla f(x)$$

$$\nabla^2 h(x) = g'(f(x))\nabla^2 f(x) + g''(f(x))\nabla f(x)\nabla f(x)^\top$$

- Composition with affine function

$$g(x) = f(Ax + b)$$

$$\nabla g(x) = A^\top \nabla f(Ax + b)$$

$$\nabla^2 g(x) = A^\top \nabla^2 f(Ax + b)A$$



# Second Derivative

## □ Chain rule

- Suppose  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g: \mathbf{R} \rightarrow \mathbf{R}$ , and  $h(x) = g(f(x))$

$$\nabla h(x) = g'(f(x))\nabla f(x)$$

$$\nabla^2 h(x) = g'(f(x))\nabla^2 f(x) + g''(f(x))\nabla f(x)\nabla f(x)^\top$$

- Composition with affine function

$$g(t) = f(x + tv), \quad x, v \in \mathbf{R}^n$$

$$g'(t) = v^\top \nabla f(x + tv)$$

$$g''(t) = v^\top \nabla^2 f(x + tv)v$$



# Example 1

---

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$g(y) = \log \sum_{i=1}^m \exp(y_i)$$

$$\nabla g(y) = \frac{1}{\sum_{i=1}^m \exp y_i} \begin{bmatrix} \exp y_1 \\ \vdots \\ \exp y_m \end{bmatrix}$$

$$\nabla^2 f(x) = A^\top \nabla^2 g(Ax + b) A$$





# Example 1

---

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$g(y) = \log \sum_{i=1}^m \exp(y_i)$$

$$\nabla g(y) = \frac{1}{\sum_{i=1}^m \exp y_i} \begin{bmatrix} \exp y_1 \\ \vdots \\ \exp y_m \end{bmatrix}$$

$$\nabla^2 g(y) = \text{diag}(\nabla g(y)) - \nabla g(y) \nabla g(y)^\top$$



# Example 1

□ Consider the function  $f: \mathbf{R}^n \rightarrow \mathbf{R}$

$$f(x) = \log \sum_{i=1}^m \exp(a_i^\top x + b_i)$$

■ where  $a_1, \dots, a_m \in \mathbf{R}^n$ ,  $b_1, \dots, b_m \in \mathbf{R}$

□  $f(x) = g(Ax + b)$

$$\begin{aligned} \nabla^2 f(x) &= A^\top \nabla g^2(Ax + b) A \\ &= A^\top \left( \frac{1}{\mathbf{1}^\top z} \text{diag}(z) - \frac{1}{(\mathbf{1}^\top z)^2} z z^\top \right) A \end{aligned}$$

■  $z_i = \exp(a_i^\top x + b_i)$ ,  $i = 1, \dots, m$



# Outline

---

- Norms
- Analysis
- Functions
- Derivatives
- Linear Algebra



# Linear algebra

---

## □ Range and nullspace

- Let  $A \in \mathbf{R}^{m \times n}$ , the range of  $A$ , denoted  $\mathcal{R}(A)$ , is the set of all vectors in  $\mathbf{R}^m$  that can be written as linear combinations of the columns of  $A$ :

$$\mathcal{R}(A) = \{Ax | x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$$

- The nullspace (or kernel) of  $A$ , denoted  $\mathcal{N}(A)$ , is the set of all vectors  $x$  mapped into zero by  $A$ :

$$\mathcal{N}(A) = \{x | Ax = 0\} \subseteq \mathbf{R}^n$$

- If  $\mathcal{V}$  is a subspace of  $\mathbf{R}^n$ , its orthogonal complement, denoted  $\mathcal{V}^\perp$ , is defined as:

$$\mathcal{V}^\perp = \{x | z^\top x = 0 \text{ for all } z \in \mathcal{V}\}$$



# Linear algebra

## □ Range and nullspace

- Let  $A \in \mathbf{R}^{m \times n}$ , the range of  $A$ , denoted  $\mathcal{R}(A)$ , is the set of all vectors in  $\mathbf{R}^m$  that can be written as linear combinations of the columns of  $A$ :

$$\mathcal{R}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$$

- The nullspace of  $A$ , denoted  $\mathcal{N}(A)$ , is the set of all vectors in  $\mathbf{R}^n$  that are mapped into zero by  $A$ :

$$\mathcal{N}(A) = \mathcal{R}(A^T)^\perp$$

$$\mathcal{N}(A) = \{x \mid Ax = 0\} \subseteq \mathbf{R}^n$$

- If  $\mathcal{V}$  is a subspace of  $\mathbf{R}^n$ , its orthogonal complement, denoted  $\mathcal{V}^\perp$ , is defined as:

$$\mathcal{V}^\perp = \{x \mid z^T x = 0 \text{ for all } z \in \mathcal{V}\}$$



# Linear algebra

---

## □ Symmetric eigenvalue decomposition

- Suppose  $A \in \mathbf{S}^n$ , i.e.,  $A$  is a real symmetric  $n \times n$  matrix. Then  $A$  can be factored as

$$A = Q\Lambda Q^T$$

where  $Q \in \mathbf{R}^{n \times n}$  is orthogonal, i.e., satisfies  $Q^T Q = I$ , and  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$

- The determinant and trace can be expressed in terms of the eigenvalue

$$\det A = \prod_{i=1}^n \lambda_i, \text{tr } A = \sum_{i=1}^n \lambda_i$$



# Linear algebra

---

## □ Norms

$$\|A\|_2 = \max_{i=1,\dots,n} |\lambda_i| = \max(\lambda_1, -\lambda_n)$$

$$\|A\|_F = \left( \sum_{i=1}^n \lambda_i^2 \right)^{1/2}$$



# Linear algebra

---

## □ Positive Definite Matrix

- A matrix  $A \in \mathbf{S}^n$  is called **positive definite**, if for all  $x \neq 0$ ,  $x^T A x > 0$ , denoted as  $A \succ 0$ .
  - ✓ If and only all eigenvalues are positive
- If  $-A$  is positive definite, we say  $A$  is negative definite, denoted as  $A \prec 0$ .
- We use  $\mathbf{S}_{++}^n$  to denote the set of positive definite matrices in  $\mathbf{S}^n$ .





# Linear algebra

---

## □ Positive Semidefinite Matrix

- A matrix  $A \in \mathbf{S}^n$  is called **positive semidefinite**, if for all  $x \neq 0, x^T A x \geq 0$ , denoted as  $A \succcurlyeq 0$ .
  - ✓ If and only all eigenvalues are nonnegative
- If  $-A$  is positive semidefinite, we say  $A$  is negative semidefinite, denoted as  $A \preccurlyeq 0$ .
- We use  $\mathbf{S}_+^n$  to denote the set of positive semidefinite matrices in  $\mathbf{S}^n$ .



# Linear algebra

---

## □ Singular value decomposition (SVD)

- Suppose  $A \in \mathbf{R}^{m \times n}$  with  $\text{rank } A = r$ . Then  $A$  can be factored as

$$A = U\Sigma V^T$$

where  $U \in \mathbf{R}^{m \times r}$  satisfies  $U^T U = I$ ,  $V \in \mathbf{R}^{n \times r}$  satisfies  $V^T V = I$ , and  $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_r)$  with

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$$

- The singular value decomposition can be written

$$A = \sum_{i=1}^r \sigma_i u_i v_i^T$$



# Linear algebra

---

## □ Norms

$$\|A\|_2 = \sigma_1$$

$$\|A\|_F = \left( \sum_{i=1}^n \sigma_i^2 \right)^{1/2}$$



# Discussions

---

## □ Symmetric eigenvalue decomposition

- Suppose  $A \in \mathbf{S}^n$

$$A = Q\Lambda Q^T$$

## □ Singular value decomposition (SVD)

- Suppose  $A \in \mathbf{S}^n$

$$A = ?$$



# Linear algebra

---

## □ Pseudo-inverse

- Let  $A = U\Sigma V^T$  be the singular value decomposition of  $A \in \mathbf{R}^{m \times n}$ , with  $\text{rank } A = r$ . The pseudo-inverse or Moore-Penrose inverse of  $A$  is

$$A^\dagger = V\Sigma^{-1}U^T \in \mathbf{R}^{n \times m}$$
$$AA^\dagger A = A$$

## □ Schur complement

- $A \in \mathbf{S}^k$ , and a matrix  $X \in \mathbf{S}^n$  partitioned as

$$X = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$$

- If  $\det A \neq 0$ , the matrix

$$S = C - B^T A^{-1} B$$

is called the Schur complement of  $A$  in  $X$

# Application of Schur complement



## □ Determinant

$$\det X = \det A \det S$$

## □ PD Matrices

- $X \succ 0$  if and only if  $A \succ 0$  and  $S \succ 0$
- If  $A \succ 0$ , then  $X \succcurlyeq 0$  if and only if  $S \succcurlyeq 0$

## □ PSD Matrices

$$X \succcurlyeq 0 \iff A \succcurlyeq 0, (I - AA^\dagger)B = 0, C - B^\top A^\dagger B \succcurlyeq 0$$



# Summary

---

- Norms of vectors
  - $l_1$ -norm,  $l_2$ -norm,  $l_\infty$ -norm,  $P$ -quadratic norm
- Norms of Matrices
  - Frobenius norm, spectral norm, nuclear norm
- Gradients of Common Functions
  - The Matrix Cookbook
- Eigendecomposition vs SVD
- PSD matrices