





Lecture 11: Uncertainty 2

http://cs.nju.edu.cn/yuy/course_ai17.ashx



Previously...



Conditional Probability
Conditional Independence

Bayesian Network:
a network of conditional independence

Constructing Bayesian networks



Need a method such that a series of locally testable assertions of conditional independence guarantees the required global semantics


1. Choose an ordering of variables X_1, \dots, X_n
2. For $i = 1$ to n
 - add X_i to the network
 - select parents from X_1, \dots, X_{i-1} such that
$$\mathbf{P}(X_i | Parents(X_i)) = \mathbf{P}(X_i | X_1, \dots, X_{i-1})$$

This choice of parents guarantees the global semantics:

$$\begin{aligned}\mathbf{P}(X_1, \dots, X_n) &= \prod_{i=1}^n \mathbf{P}(X_i | X_1, \dots, X_{i-1}) \quad (\text{chain rule}) \\ &= \prod_{i=1}^n \mathbf{P}(X_i | Parents(X_i)) \quad (\text{by construction})\end{aligned}$$

Example

Suppose we choose the ordering M, J, A, B, E



$P(J|M) = P(J)$? No

$P(A|J, M) = P(A|J)$? $P(A|J, M) = P(A)$? No

$P(B|A, J, M) = P(B|A)$? Yes

$P(B|A, J, M) = P(B)$? No

$P(E|B, A, J, M) = P(E|A)$? No

$P(E|B, A, J, M) = P(E|A, B)$? Yes




Example: Car diagnosis



Initial evidence: car won't start

Testable variables (green), "broken, so fix it" variables (orange)

Hidden variables (gray) ensure sparse structure, reduce parameters



Compact conditional distributions



CPT grows exponentially with number of parents

CPT becomes infinite with continuous-valued parent or child

Solution: **canonical** distributions that are defined compactly

Deterministic nodes are the simplest case:

$$X = f(\text{Parents}(X)) \text{ for some function } f$$

E.g., Boolean functions

$$\text{NorthAmerican} \Leftrightarrow \text{Canadian} \vee \text{US} \vee \text{Mexican}$$

E.g., numerical relationships among continuous variables

$$\frac{\partial \text{Level}}{\partial t} = \text{inflow} + \text{precipitation} - \text{outflow} - \text{evaporation}$$

Compact conditional distributions contd.



Noisy-OR distributions model multiple noninteracting causes


- 1) Parents $U_1 \dots U_k$ include all causes (can add **leak node**)
- 2) Independent failure probability q_i for each cause alone

$$\Rightarrow P(X|U_1 \dots U_j, \neg U_{j+1} \dots \neg U_k) = 1 - \prod_{i=1}^j q_i$$


<i>Cold</i>	<i>Flu</i>	<i>Malaria</i>	$P(\text{Fever})$	$P(\neg \text{Fever})$
F	F	F	0.0	1.0
F	F	T	0.9	0.1
F	T	F	0.8	0.2
F	T	T	0.98	$0.02 = 0.2 \times 0.1$
T	F	F	0.4	0.6
T	F	T	0.94	$0.06 = 0.6 \times 0.1$
T	T	F	0.88	$0.12 = 0.6 \times 0.2$
T	T	T	0.988	$0.012 = 0.6 \times 0.2 \times 0.1$

Number of parameters **linear** in number of parents

Hybrid (discrete+continuous) networks



Discrete (*Subsidy?* and *Buys?*); continuous (*Harvest* and *Cost*)



Option 1: discretization—possibly large errors, large CPTs

Option 2: finitely parameterized canonical families

- 1) Continuous variable, discrete+continuous parents (e.g., *Cost*)
- 2) Discrete variable, continuous parents (e.g., *Buys?*)



Continuous child variables

Need one **conditional density** function for child variable given continuous parents, for each possible assignment to discrete parents

Most common is the **linear Gaussian** model, e.g.,:


$$\begin{aligned} P(\textit{Cost} = c | \textit{Harvest} = h, \textit{Subsidy?} = \textit{true}) \\ &= N(a_t h + b_t, \sigma_t)(c) \\ &= \frac{1}{\sigma_t \sqrt{2\pi}} \exp\left(-\frac{1}{2} \left(\frac{c - (a_t h + b_t)}{\sigma_t}\right)^2\right) \end{aligned}$$

Mean *Cost* varies linearly with *Harvest*, variance is fixed

Linear variation is unreasonable over the full range

but works OK if the **likely** range of *Harvest* is narrow

Continuous child variables




All-continuous network with LG distributions


⇒ full joint distribution is a multivariate Gaussian

Discrete+continuous LG network is a **conditional Gaussian** network i.e., a multivariate Gaussian over all continuous variables for each combination of discrete variable values

Discrete variable w/ continuous parents



Probability of *Buys?* given *Cost* should be a “soft” threshold:



Probit distribution uses integral of Gaussian:


$$\Phi(x) = \int_{-\infty}^x N(0, 1)(x)dx$$

$$P(\text{Buys?} = \text{true} \mid \text{Cost} = c) = \Phi((-c + \mu)/\sigma)$$


Why the probit?



1. It's sort of the right shape
2. Can view as hard threshold whose location is subject to noise




Discrete variable contd.




Sigmoid (or logit) distribution also used in neural networks:

$$P(\text{Buys?} = \text{true} \mid \text{Cost} = c) = \frac{1}{1 + \exp\left(-2\frac{-c+\mu}{\sigma}\right)}$$

Sigmoid has similar shape to probit but much longer tails:





Inference in Bayesian networks

Inference tasks



Simple queries: compute posterior marginal $\mathbf{P}(X_i|\mathbf{E} = \mathbf{e})$

e.g., $P(\text{NoGas}|\text{Gauge} = \text{empty}, \text{Lights} = \text{on}, \text{Starts} = \text{false})$

Conjunctive queries: $\mathbf{P}(X_i, X_j|\mathbf{E} = \mathbf{e}) = \mathbf{P}(X_i|\mathbf{E} = \mathbf{e})\mathbf{P}(X_j|X_i, \mathbf{E} = \mathbf{e})$

Optimal decisions: decision networks include utility information;
probabilistic inference required for $P(\text{outcome}|\text{action}, \text{evidence})$

Value of information: which evidence to seek next?

Sensitivity analysis: which probability values are most critical?

Explanation: why do I need a new starter motor?

Exact inference




Inference by enumeration

Slightly intelligent way to sum out variables from the joint without actually constructing its explicit representation

Simple query on the burglary network:

$$\begin{aligned} & \mathbf{P}(B|j, m) \\ &= \mathbf{P}(B, j, m) / P(j, m) \\ &= \alpha \mathbf{P}(B, j, m) \\ &= \alpha \sum_e \sum_a \mathbf{P}(B, e, a, j, m) \end{aligned}$$




Rewrite full joint entries using product of CPT entries:

$$\begin{aligned} & \mathbf{P}(B|j, m) \\ &= \alpha \sum_e \sum_a \mathbf{P}(B)P(e)\mathbf{P}(a|B, e)P(j|a)P(m|a) \\ &= \alpha \mathbf{P}(B) \sum_e P(e) \sum_a \mathbf{P}(a|B, e)P(j|a)P(m|a) \end{aligned}$$

Recursive depth-first enumeration: $O(n)$ space, $O(d^n)$ time

Enumeration algorithm



function **ENUMERATION-ASK**(X, \mathbf{e}, bn) **returns** a distribution over X

inputs: X , the query variable

\mathbf{e} , observed values for variables \mathbf{E}

bn , a Bayesian network with variables $\{X\} \cup \mathbf{E} \cup \mathbf{Y}$

$Q(X) \leftarrow$ a distribution over X , initially empty

for each value x_i of X **do**

 extend \mathbf{e} with value x_i for X

$Q(x_i) \leftarrow$ **ENUMERATE-ALL**(**VARS**[bn], \mathbf{e})

return **NORMALIZE**($Q(X)$)

function **ENUMERATE-ALL**($vars, \mathbf{e}$) **returns** a real number

if **EMPTY?**($vars$) **then return** 1.0

$Y \leftarrow$ **FIRST**($vars$)


if Y has value y in \mathbf{e}

then return $P(y \mid Pa(Y)) \times$ **ENUMERATE-ALL**(**REST**($vars$), \mathbf{e})

else return $\sum_y P(y \mid Pa(Y)) \times$ **ENUMERATE-ALL**(**REST**($vars$), \mathbf{e}_y)

 where \mathbf{e}_y is \mathbf{e} extended with $Y = y$

Evaluation tree



Enumeration is inefficient: repeated computation

e.g., computes $P(j|a)P(m|a)$ for each value of e

Inference by variable elimination



Variable elimination: carry out summations right-to-left, storing intermediate results (**factors**) to avoid recomputation

$$\begin{aligned} \mathbf{P}(B|j, m) &= \alpha \underbrace{\mathbf{P}(B)}_B \underbrace{\sum_e P(e)}_E \underbrace{\sum_a \mathbf{P}(a|B, e)}_A \underbrace{P(j|a)}_J \underbrace{P(m|a)}_M \\ &= \alpha \mathbf{P}(B) \sum_e P(e) \sum_a \mathbf{P}(a|B, e) P(j|a) f_M(a) \\ &= \alpha \mathbf{P}(B) \sum_e P(e) \sum_a \mathbf{P}(a|B, e) f_J(a) f_M(a) \\ &= \alpha \mathbf{P}(B) \sum_e P(e) \sum_a f_A(a, b, e) f_J(a) f_M(a) \\ &= \alpha \mathbf{P}(B) \sum_e P(e) f_{\bar{A}JM}(b, e) \text{ (sum out } A) \\ &= \alpha \mathbf{P}(B) f_{\bar{E}\bar{A}JM}(b) \text{ (sum out } E) \\ &= \alpha f_B(b) \times f_{\bar{E}\bar{A}JM}(b) \end{aligned}$$

Variable elimination: Basic operations



Summing out a variable from a product of factors:

move any constant factors outside the summation

add up submatrices in pointwise product of remaining factors

$$\sum_x f_1 \times \cdots \times f_k = f_1 \times \cdots \times f_i \sum_x f_{i+1} \times \cdots \times f_k = f_1 \times \cdots \times f_i \times f_{\bar{X}}$$


assuming f_1, \dots, f_i do not depend on X

Pointwise product of factors f_1 and f_2 :

$$\begin{aligned} f_1(x_1, \dots, x_j, y_1, \dots, y_k) \times f_2(y_1, \dots, y_k, z_1, \dots, z_l) \\ = f(x_1, \dots, x_j, y_1, \dots, y_k, z_1, \dots, z_l) \end{aligned}$$

E.g., $f_1(a, b) \times f_2(b, c) = f(a, b, c)$

Variable elimination algorithm



```
function ELIMINATION-ASK( $X, \mathbf{e}, bn$ ) returns a distribution over  $X$   
inputs:  $X$ , the query variable  
           $\mathbf{e}$ , evidence specified as an event  
           $bn$ , a belief network specifying joint distribution  $\mathbf{P}(X_1, \dots, X_n)$   
 $factors \leftarrow []$ ;  $vars \leftarrow \text{REVERSE}(\text{VARS}[bn])$   
for each  $var$  in  $vars$  do  
     $factors \leftarrow [\text{MAKE-FACTOR}(var, \mathbf{e}) | factors]$   
    if  $var$  is a hidden variable then  $factors \leftarrow \text{SUM-OUT}(var, factors)$   
return  $\text{NORMALIZE}(\text{POINTWISE-PRODUCT}(factors))$ 
```


Irrelevant variables



Consider the query $P(\text{JohnCalls} | \text{Burglary} = \text{true})$

$$P(J|b) = \alpha P(b) \sum_e P(e) \sum_a P(a|b, e) P(J|a) \sum_m P(m|a)$$

Sum over m is identically 1; M is **irrelevant** to the query



Thm 1: Y is irrelevant unless $Y \in \text{Ancestors}(\{X\} \cup \mathbf{E})$

Here, $X = \text{JohnCalls}$, $\mathbf{E} = \{\text{Burglary}\}$, and
 $\text{Ancestors}(\{X\} \cup \mathbf{E}) = \{\text{Alarm}, \text{Earthquake}\}$
so MaryCalls is irrelevant

(Compare this to backward chaining from the query in Horn clause KBs)

Irrelevant variables contd.





Defn: moral graph of Bayes net: marry all parents and drop arrows

Defn: **A** is m-separated from **B** by **C** iff separated by **C** in the moral graph

Thm 2: **Y** is irrelevant if m-separated from **X** by **E**

For $P(\text{JohnCalls} | \text{Alarm} = \text{true})$, both *Burglary* and *Earthquake* are irrelevant





Complexity of exact inference


Singly connected networks (or **polytrees**):

- any two nodes are connected by at most one (undirected) path
- time and space cost of variable elimination are $O(d^k n)$


Multiply connected networks:

- can reduce 3SAT to exact inference \Rightarrow NP-hard
- equivalent to **counting** 3SAT models \Rightarrow #P-complete

1. $A \vee B \vee C$
2. $C \vee D \vee \neg A$
3. $B \vee C \vee \neg D$



Approximate inference



Inference by stochastic simulation

Basic idea:

- 1) Draw N samples from a sampling distribution S
- 2) Compute an approximate posterior probability \hat{P}
- 3) Show this converges to the true probability P

Outline:

- Sampling from an empty network
- Rejection sampling: reject samples disagreeing with evidence
- Likelihood weighting: use evidence to weight samples
- Markov chain Monte Carlo (MCMC): sample from a stochastic process whose stationary distribution is the true posterior



About random number generation



How to generate a discrete distribution from the uniform distribution?

given $U[0,1]$

generate A 30%, B 60%, C 10%

About random number generation




How to generate a continuous distribution from the uniform distribution?

given $U[0,1]$

generate $N(0,1)$

About random number generation




How to generate a discrete distribution from a discrete distribution?


given A,B,C 33.33%

generate A,B,C,D 25%


Sampling from an empty network




Sampling from an empty network




Sampling from an empty network




Sampling from an empty network




Sampling from an empty network



Sampling from an empty network



Sampling from an empty network




Sampling from an empty network



```
function PRIOR-SAMPLE(bn) returns an event sampled from bn  
  inputs: bn, a belief network specifying joint distribution  $\mathbf{P}(X_1, \dots, X_n)$   
  x  $\leftarrow$  an event with n elements  
  for i = 1 to n do  
    xi  $\leftarrow$  a random sample from  $\mathbf{P}(X_i \mid \text{parents}(X_i))$   
      given the values of Parents(Xi) in x  
  return x
```

Sampling from an empty network contd.



Probability that PRIORSAMPLE generates a particular event

$$S_{PS}(x_1 \dots x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i)) = P(x_1 \dots x_n)$$

i.e., the true prior probability

E.g., $S_{PS}(t, f, t, t) = 0.5 \times 0.9 \times 0.8 \times 0.9 = 0.324 = P(t, f, t, t)$

Let $N_{PS}(x_1 \dots x_n)$ be the number of samples generated for event x_1, \dots, x_n


Then we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \hat{P}(x_1, \dots, x_n) &= \lim_{N \rightarrow \infty} N_{PS}(x_1, \dots, x_n) / N \\ &= S_{PS}(x_1, \dots, x_n) \\ &= P(x_1 \dots x_n) \end{aligned}$$

That is, estimates derived from PRIORSAMPLE are **consistent**

Shorthand: $\hat{P}(x_1, \dots, x_n) \approx P(x_1 \dots x_n)$

Conditional Probability: Rejection sampling



$\hat{P}(X|e)$ estimated from samples agreeing with e

```
function REJECTION-SAMPLING( $X, e, bn, N$ ) returns an estimate of  $P(X|e)$ 
  local variables:  $N$ , a vector of counts over  $X$ , initially zero
  for  $j = 1$  to  $N$  do
     $x \leftarrow$  PRIOR-SAMPLE( $bn$ )
    if  $x$  is consistent with  $e$  then
       $N[x] \leftarrow N[x] + 1$  where  $x$  is the value of  $X$  in  $x$ 
  return NORMALIZE( $N[X]$ )
```

E.g., estimate $P(Rain|Sprinkler = true)$ using 100 samples

27 samples have $Sprinkler = true$

Of these, 8 have $Rain = true$ and 19 have $Rain = false$.

$\hat{P}(Rain|Sprinkler = true) = \text{NORMALIZE}(\langle 8, 19 \rangle) = \langle 0.296, 0.704 \rangle$

Similar to a basic real-world empirical estimation procedure

Analysis of rejection sampling



$$\begin{aligned}\hat{\mathbf{P}}(X|\mathbf{e}) &= \alpha \mathbf{N}_{PS}(X, \mathbf{e}) && \text{(algorithm defn.)} \\ &= \mathbf{N}_{PS}(X, \mathbf{e}) / N_{PS}(\mathbf{e}) && \text{(normalized by } N_{PS}(\mathbf{e})\text{)} \\ &\approx \mathbf{P}(X, \mathbf{e}) / P(\mathbf{e}) && \text{(property of PRIORSAMPLE)} \\ &= \mathbf{P}(X|\mathbf{e}) && \text{(defn. of conditional probability)}\end{aligned}$$

Hence rejection sampling returns consistent posterior estimates

Problem: hopelessly expensive if $P(\mathbf{e})$ is small

$P(\mathbf{e})$ drops off exponentially with number of evidence variables!

Likelihood weighting




Idea: fix evidence variables, sample only nonevidence variables, and weight each sample by the likelihood it accords the evidence

```
function LIKELIHOOD-WEIGHTING( $X, \mathbf{e}, bn, N$ ) returns an estimate of  $P(X|\mathbf{e})$   
  local variables:  $\mathbf{W}$ , a vector of weighted counts over  $X$ , initially zero  
  for  $j = 1$  to  $N$  do  
     $\mathbf{x}, w \leftarrow$  WEIGHTED-SAMPLE( $bn$ )  
     $\mathbf{W}[x] \leftarrow \mathbf{W}[x] + w$  where  $x$  is the value of  $X$  in  $\mathbf{x}$   
  return NORMALIZE( $\mathbf{W}[X]$ )
```


```
function WEIGHTED-SAMPLE( $bn, \mathbf{e}$ ) returns an event and a weight  
   $\mathbf{x} \leftarrow$  an event with  $n$  elements;  $w \leftarrow 1$   
  for  $i = 1$  to  $n$  do  
    if  $X_i$  has a value  $x_i$  in  $\mathbf{e}$   
      then  $w \leftarrow w \times P(X_i = x_i \mid \text{parents}(X_i))$   
      else  $x_i \leftarrow$  a random sample from  $\mathbf{P}(X_i \mid \text{parents}(X_i))$   
  return  $\mathbf{x}, w$ 
```

Likelihood weighting example




$$w = 1.0$$

Likelihood weighting example




$$w = 1.0$$

Likelihood weighting example




$$w = 1.0$$

Likelihood weighting example




$$w = 1.0 \times 0.1$$

Likelihood weighting example



$$w = 1.0 \times 0.1$$

Likelihood weighting example



$$w = 1.0 \times 0.1 \times 0.99 = 0.099$$

Likelihood weighting analysis



Sampling probability for WEIGHTEDSAMPLE is

$$S_{WS}(\mathbf{z}, \mathbf{e}) = \prod_{i=1}^l P(z_i | \text{parents}(Z_i))$$

Note: pays attention to evidence in **ancestors** only


⇒ somewhere “in between” prior and posterior distribution

Weight for a given sample \mathbf{z}, \mathbf{e} is

$$w(\mathbf{z}, \mathbf{e}) = \prod_{i=1}^m P(e_i | \text{parents}(E_i))$$

Weighted sampling probability is

$$\begin{aligned} & S_{WS}(\mathbf{z}, \mathbf{e})w(\mathbf{z}, \mathbf{e}) \\ &= \prod_{i=1}^l P(z_i | \text{parents}(Z_i)) \prod_{i=1}^m P(e_i | \text{parents}(E_i)) \\ &= P(\mathbf{z}, \mathbf{e}) \text{ (by standard global semantics of network)} \end{aligned}$$



Hence likelihood weighting returns consistent estimates but performance still degrades with many evidence variables because a few samples have nearly all the total weight

Approximate inference using MCMC



“State” of network = current assignment to all variables.

Generate next state by sampling one variable given Markov blanket

Sample each variable in turn, keeping evidence fixed


```
function MCMC-Ask( $X, \mathbf{e}, bn, N$ ) returns an estimate of  $P(X|\mathbf{e})$   
  local variables:  $\mathbf{N}[X]$ , a vector of counts over  $X$ , initially zero  
     $\mathbf{Z}$ , the nonevidence variables in  $bn$   
     $\mathbf{x}$ , the current state of the network, initially copied from  $\mathbf{e}$   
  
  initialize  $\mathbf{x}$  with random values for the variables in  $\mathbf{Y}$   
  for  $j = 1$  to  $N$  do  
    for each  $Z_i$  in  $\mathbf{Z}$  do  
      sample the value of  $Z_i$  in  $\mathbf{x}$  from  $\mathbf{P}(Z_i|mb(Z_i))$   
        given the values of  $MB(Z_i)$  in  $\mathbf{x}$   
       $\mathbf{N}[x] \leftarrow \mathbf{N}[x] + 1$  where  $x$  is the value of  $X$  in  $\mathbf{x}$   
  return NORMALIZE( $\mathbf{N}[X]$ )
```

Can also choose a variable to sample at random each time

The Markov chain



With $Sprinkler = true$, $WetGrass = true$, there are four states:



Wander about for a while, average what you see

MCMC example contd.



Estimate $\mathbf{P}(Rain|Sprinkler = true, WetGrass = true)$

Sample *Cloudy* or *Rain* given its Markov blanket, repeat.
Count number of times *Rain* is true and false in the samples.

E.g., visit 100 states

31 have *Rain = true*, 69 have *Rain = false*

$$\begin{aligned} \hat{\mathbf{P}}(Rain|Sprinkler = true, WetGrass = true) \\ = \text{NORMALIZE}(\langle 31, 69 \rangle) = \langle 0.31, 0.69 \rangle \end{aligned}$$

Theorem: chain approaches **stationary distribution**:

long-run fraction of time spent in each state is exactly proportional to its posterior probability

Markov blanket sampling




Markov blanket of *Cloudy* is

Sprinkler and *Rain*

Markov blanket of *Rain* is

Cloudy, *Sprinkler*, and *WetGrass*



Probability given the Markov blanket is calculated as follows:

$$P(x'_i | mb(X_i)) = P(x'_i | parents(X_i)) \prod_{Z_j \in Children(X_i)} P(z_j | parents(Z_j))$$

Easily implemented in message-passing parallel systems, brains

Main computational problems:

- 1) Difficult to tell if convergence has been achieved
- 2) Can be wasteful if Markov blanket is large:

$P(X_i | mb(X_i))$ won't change much (law of large numbers)

Summary



Exact inference by variable elimination:

- polytime on polytrees, NP-hard on general graphs
- space = time, very sensitive to topology

Approximate inference by LW, MCMC:

- LW does poorly when there is lots of (downstream) evidence
- LW, MCMC generally insensitive to topology
- Convergence can be very slow with probabilities close to 1 or 0
- Can handle arbitrary combinations of discrete and continuous variables